


Aspergillosis And The Lungs

Fungal Disease Series

Aspergillosis (As-per-gill-osis) is an infection caused by a fungus called *Aspergillus*. *Aspergillus* lives in soil, plants and rotting material. It can also be found in the dust in your home, carpeting, heating and air conditioning ducts, certain foods including dried fish and in marijuana. Aspergillus infection is occurring more often and is now the leading cause of death due to invasive fungal infections in the United States. This is due mainly because there are more people who are living with weakened immune systems that put them at higher risk of infection.



Not everyone who gets aspergillosis goes on to develop the severe form (*invasive aspergillosis*).

What causes Aspergillosis?

Aspergillus enters the body when you breathe in the fungal spores (“seeds”). This fungus is commonly found in your lungs and sinuses. If your immunity (the ability to “fight off” infections) is normal, the infection can be contained and may never cause an illness. However, having a weak immune system or a chronic lung disease allows the *Aspergillus* to grow, invade your lungs and spread throughout your body. This may happen if you:

- have a cancer such as leukemia or aplastic anemia;
- take chemotherapy or are on corticosteroids for a long time (for any reason);
- had an organ transplant (for example, kidney or lung);
- have advanced HIV infection; or
- have a chronic lung disease like asthma, emphysema, tuberculosis, or cystic fibrosis.

What are the different forms and symptoms of aspergillosis?

There are several different ways a person can react to this fungus, depending on one’s immunity and other

factors. Different forms and their symptoms include:

- **Hypersensitivity Pneumonitis**—an allergic reaction to the fungus in the lungs. Symptoms can last for weeks or months and include:
 - shortness of breath
 - coughing
- **Allergic Bronchopulmonary Aspergillosis (ABPA)**—an asthma-like illness. Symptoms do not improve with usual asthma treatment and include:
 - coughing
 - shortness of breath
 - wheezing
- **Invasive Aspergillosis**—a rapidly spreading and potentially life threatening illness. Symptoms include:
 - fever that does not get better with antibiotics
 - difficulty breathing
 - chest pain that is worse with deep breathing
 - dry cough
 - coughing up blood, which can be caused by a ball of fungus that forms in the lungs (called an aspergilloma)

In severe cases of invasive disease, additional symptoms include:

- Sinus infection leading to swollen eye on one side or bleeding from nose

- difficulty talking
- paralysis of facial muscles
- ulcers inside the mouth or inside the chest wall
- confusion, seizures or stroke-like symptoms, which could mean the infection has spread to the brain.

How is Aspergillosis diagnosed?

Aspergillosis generally starts out as a spot in your lung called a lung “nodule”. (This nodule can be mistaken for lung cancer or tuberculosis). When the fungus is just a nodule, you likely will not have symptoms. However, over time, the fungus may develop into pulmonary nodules, a fungus ball, or pneumonia. (see ATS handouts on Pulmonary Nodules and Pneumonia at www.thoracic.org/patients/).

Unfortunately, the only way to diagnose aspergillosis with certainty is by a tissue biopsy (sampling a piece of your lung). Other tests that can be done include examining your sputum (mucus or phlegm) or taking cultures from your airways (breathing tubes). Culture specimens can be collected by a bronchoscopy (see ATS handout on Flexible Bronchoscopy at www.thoracic.org/patients/). Unfortunately, these cultures sometimes miss the infection. There is also a test that identifies parts of the fungus in your blood, called the *serum galactomannan test*. However, this test is not 100% accurate. It may miss a few cases, but a positive test increases the chance you have the infection, especially if you have risk factors for aspergillosis, and have what looks like aspergillosis on a chest X-ray or CT scan.

How is aspergillosis treated?

If you develop symptoms of aspergillosis, you may need treatment with an anti-fungal drug for either a brief or a long time. The worse your symptoms are, the more aggressive your healthcare provider will be in diagnosing and starting treatment. Anti-fungal drugs used to treat aspergillosis include: voriconazole, amphotericin B, caspofungin, itraconazole, and posaconazole. Because most of these drugs have side effects, they are not usually given unless your healthcare provider is sure that you have aspergillosis. If you have a history of aspergillosis, you may be asked to take an anti-fungal drug to prevent the infection from coming back, especially when your immune system is weak (such as if you are receiving a course of cancer chemotherapy). If you have an *aspergilloma*, you may need surgery to remove the fungus ball because drugs are not very effective in treating it. Surgery may also be recommended if your infection does not improve with drugs.

Treatment of allergic bronchopulmonary *aspergillosis* (ABPA) is aimed at preventing and treating flare-ups and preventing damage to your lungs and other organs. Treatment for allergic reactions includes a corticosteroid by mouth and an anti-fungal drug like itraconazole, which are often taken for several months.

Can aspergillosis be prevented?

It is difficult to avoid being exposed to *Aspergillus*. However, if your immune system is weak, you should take special precautions. For example, avoid dusty areas, or at least wear an N95 mask (see ATS handout on Disposable Respirators at www.thoracic.org/patients/), when you are around dusty areas, and avoid gardening until your immune system improves.

Use of anti-fungal drugs to prevent aspergillosis is helpful, mainly if you are at high-risk (have a low white blood cell count, leukemia or a myelodysplastic syndrome, or if you have received a stem cell transplant). Your healthcare provider will advise you if you need preventive treatment.

Author: Hrishikesh S Kulkarni MD

Reviewers: Suzanne Lareau RN, MS, Bonnie Fahy RN, MN, Chadi Hage MD, Marianna Sockrider MD, DrPH

Rx Action Steps

See your healthcare provider if you notice any of these symptoms:

- ✓ A fever that will not go away, especially if you are also having shortness of breath and/or chest pain, and if your immunity is weak
- ✓ Chronic cough, especially if your sputum (phlegm) is blood-stained
- ✓ Frequent asthma flares that do not improve even when you take your usual asthma medicines

Doctor's Office Telephone:

Resources:

Centers for Disease Control and Prevention (CDC)
<https://www.cdc.gov/fungal/diseases/aspergillosis/>

Support for People Living with Aspergillosis
<http://www.nacpatients.org.uk/>

Medline Plus
<https://medlineplus.gov/aspergillosis.html>

This information is a public service of the American Thoracic Society. The content is for educational purposes only. It should not be used as a substitute for the medical advice of one's healthcare provider.

